

EDITORIAL COMMENT

Botulinum Toxin Type A as a Prophylactic Treatment for Chronic Daily Headache: Time to Conduct a Large Clinical Trial in Taiwan?

Hsiu-Chih Liu*

Department of Neurology, National Yang-Ming University School of Medicine and Neurological Institute, Taipei Veterans General Hospital, Taipei, Taiwan, R.O.C.

Chronic daily headache (CDH) refers to a group of headache disorders that occur on at least 15 days per month.¹ It is frequently associated with the overuse of pain medications and comorbid psychiatric conditions such as anxiety and depression. Although CDH is not life-threatening, it often causes considerable disability and compromises the quality of life of those who suffer from it. Therefore, the management of CDH is a challenge for many clinicians.²

One of the most common subtypes of CDH is transformed migraine (TM), also known as chronic migraine (CM). Patients with TM usually have a history of episodic migraine which over the years gradually increases in frequency to become CDH, but decrease in the severity of migraine features.¹ In TM, acute migraine attacks are treated as migraine, but their prevention is difficult.³ Conventional preventive treatments for TM include the use of β -blockers, tricyclic antidepressants, and anticonvulsants, but the results are less than satisfactory.² The systemic side effects of these medications also limit their usefulness as prophylactic treatments. To date, there are no randomized controlled trials of preventive therapy aimed exclusively at TM.

In the past 10 years, botulinum toxin A (BoNT-A) has emerged as a treatment for headaches, and it has been found to be safe and well tolerated in general.⁴ Several open-label studies of BoNT-A treatment with small numbers of patients have reported that it is beneficial for CDH. In this issue, Liu et al⁵ retrospectively analyzed the efficacy of BoNT-A in 30 consecutive patients with TM. After a single injection of a low dose of BoNT-A, 27 of the 30 patients reported at least

a 50% reduction in either the number of headache days or in headache intensity. The authors concluded that BoNT-A is a promising prophylactic treatment for TM for patients in Taiwan.

On the other hand, 3 randomized, double-blind, placebo-controlled studies of BoNT-A for the treatment of CDH found negative results for their primary efficacy endpoint measured by the change from baseline in the frequency of headache-free days.⁶⁻⁸ All 3 studies included patients with TM, but did not separately examine the efficacy of BoNT-A for those patients. Subgroup analysis of the study by Mathew et al⁸ showed a significant reduction in the frequency of headaches in CDH patients who were not using other prophylactic medications.⁹

The mixed findings of the efficacy of BoNT-A in CDH prophylaxis shows the need to identify those patients who may benefit from BoNT-A. The report by Liu et al⁵ indicates that it is likely that patients with TM may represent a subgroup of migrainers who respond better to BoNT-A, but further studies are needed. This issue is important in clinical practice because BoNT-A is costly and at the present time its use for headache prevention is not reimbursed by the National Health Insurance of Taiwan.

In the past decade, the epidemiology of headache and its subtypes has been extensively studied. Wang¹⁰ reported that the prevalence rates of CDH in Taiwan (3.2–3.9%) were similar to those in Western countries (3.0–4.7%). However, the inclination to seek medical attention and the response to BoNT-A treatment might be different between the 2 groups. Perhaps it is

*Correspondence to: Dr Hsiu-Chih Liu, Neurological Institute, Taipei Veterans General Hospital, 201, Section 2, Shih-Pai Road, Taipei 112, Taiwan, R.O.C.

E-mail: hcliu@vghtpe.gov.tw • Received: October 22, 2007 • Accepted: October 25, 2007

time to conduct a well-designed, large, multicenter, randomized, double-blind, placebo-controlled study of BoNT-A for the prevention of CDH in Taiwan.

References

1. Silberstein SD, Lipton RB. Chronic daily headache, including transformed migraine, chronic tension-type headache, and medication overuse. In: Silberstein SD, Lipton RB, Dalessio DJ, eds. *Wolff's Headache and Other Head Pain*. New York: Oxford University Press, 2001:247–82.
2. Saper JR, Dodick D, Gladstone JP. Management of chronic daily headache: challenges in clinical practice. *Headache* 2005; 45(Suppl):S74–85.
3. Mathew NT. The prophylactic treatment of chronic daily headache. *Headache* 2006;46:1552–64.
4. Evers S. Status on the use of botulism toxin for headache disorders. *Curr Opin Neurol* 2006;19:310–5.
5. Liu YC, Fuh JL, Chen RC, Lin KP, Wang SJ. Botulinum toxin type A in the prophylactic treatment of transformed migraine in Taiwanese patients: a review of 30 consecutive cases. *J Chin Med Assoc* 2007;70:535–40.
6. Ondo WG, Vuong KD, Derman HS. Botulinum toxin A for chronic daily headache: a randomized, placebo-controlled, parallel design study. *Cephalalgia* 2004;24:60–5.
7. Silberstein SD, Stark SR, Lucas SM, Christie SN, DeGryse RE, Turkel CC, for the BoNTA-039 Study Group. Botulinum toxin A for the prophylactic treatment of chronic daily headache: a randomized, double-blind, placebo-controlled trial. *Mayo Clin Proc* 2005;80:1126–37.
8. Mathew NT, Frishberg BM, Gawel M, Dimitrova R, Gibson J, Turkel C, BOTOX CDH Study Group. Botulinum toxin type A (BOTOX) for the prophylactic treatment of chronic daily headache: a randomized, double-blind, placebo-controlled trial. *Headache* 2005;45:293–307.
9. Dodick DW, Mauskop A, Elkind AH, DeGryse R, Brin MF, Silberstein SD, BOTOX CDH Study Group. Botulinum toxin type A for the prophylaxis of chronic daily headache: subgroup analysis of patients not receiving other prophylactic medications: a randomized double-blind, placebo-controlled study. *Headache* 2005;45:315–24.
10. Wang SJ. Epidemiology of migraine and other types of headache in Asia. *Curr Neurol Neurosci Rep* 2003;3:104–8.